



Instructions for Catalog # 775QR  
WatR™ Pollution Silica  
Revision 040307

Description:

- This standard is packaged in a 60 mL polyethylene bottle containing approximately 60 mL of standard concentrate.
- This concentrate is not preserved.
- The concentrate can be stored at room temperature.
- The diluted standard will contain Silica in the range of 50 to 250 mg/L.

Before you begin:

- This standard has been prepared as a concentrate and must be diluted prior to analysis.
- Be sure to report your Silica result as SiO<sub>2</sub> (not as Si).
- This standard should be analyzed as soon as possible after the concentrate is diluted.

Instructions:

1. Add 100-200 mL of deionized water to a clean 1000 mL class A volumetric flask.
2. Shake the Silica bottle prior to opening.
3. Using a clean, dry, class A pipet or flask, volumetrically transfer 50 mL of the concentrate into the 1000 mL volumetric flask.
4. Dilute the flask to final volume with deionized water.
5. Cap the flask and mix well.
6. Immediately analyze the diluted sample by your normal procedures.
7. Report your results as mg/L for the diluted sample.

Safety:

ERA products may be hazardous and are intended for use by professional laboratory personnel trained in the competent handling of such materials. Responsibility for the safe use of these products rests entirely with the buyer and/or user. Material Safety Data Sheets (MSDS) for all ERA products are available by calling 1-800-372-0122.